

Bell's Inequality

Jeffrey Epstein

March 11, 2018

These notes follow Bell's original papers on the subject.

Theorem 1. *A local hidden variable theory (a hidden variable theory in which the setting of one measurement apparatus does not affect the outcome of another) must obey the inequality*

$$|(ab)_\sigma - (bc)_\sigma| \leq 1 + (ac)_\sigma. \quad (1)$$

where $(ab)_\sigma = \langle A(a)B(b) \rangle_\sigma$ is the correlation between two ± 1 -valued random variables $A_\sigma(a)$ and $B_\sigma(b)$ that are perfectly anticorrelated when $a = b$.

Proof. Suppose we have a theory for describing a particular kind of thing. In our theory, these things are described by some object we'll call a preparation. I'm going to give you a bunch of things that are "the same" in the sense that they are all described by the same preparation π . You can perform an operation on these things that results in two ± 1 -valued random variables $A_\pi(a)$ and $B_\pi(b)$, where a and b are settings you are free to choose. If you wanted, you could calculate the correlation between these variables:

$$\langle A(a)B(b) \rangle_\pi. \quad (2)$$

We'd like to design a deterministic hidden-variable theory to reproduce the results of our theory. By this we mean that there is a "hidden variable" λ such that the values of the random variables are determined by λ . The preparations π determine probability distributions $\rho_\pi(\lambda)$, so that we may write the correlations as follows:

$$\langle A(a)B(b) \rangle_\pi = \int \rho_\pi(\lambda) A(a, \lambda) B(b, \lambda) d\lambda. \quad (3)$$

Suppose that the things I give you are described by a preparation σ such that the random variables $A_\sigma(a)$ and $B_\sigma(b)$ are perfectly anti-correlated when $a = b$:

$$\langle A(a)B(a) \rangle_\sigma = -1 \iff A_\sigma(a) = -B_\sigma(a). \quad (4)$$

Then for arbitrary settings a , b , and c , we can do some calculations:

$$\langle A(b)B(a) \rangle_\sigma - \langle A(b)B(c) \rangle_\sigma = \int \rho_\sigma(\lambda) [A(b, \lambda)B(a, \lambda) - A(b, \lambda)B(c, \lambda)] d\lambda \quad (5)$$

$$= \int \rho_\sigma(\lambda) [A(b, \lambda)A(c, \lambda) - A(b, \lambda)A(a, \lambda)] d\lambda \quad (6)$$

$$= \int \rho_\sigma(\lambda) A(b, \lambda) A(a, \lambda) [A(a, \lambda)A(c, \lambda) - 1] d\lambda. \quad (7)$$

Taking the absolute value and using the fact that the random variables take values ± 1 , we have

$$|\langle A(b)B(a) \rangle_\sigma - \langle A(b)B(c) \rangle_\sigma| \leq \int \rho_\sigma(\lambda) [1 - A(a, \lambda)A(c, \lambda)] d\lambda \quad (8)$$

$$= 1 + \int \rho_\sigma(\lambda) A(a, \lambda) B(c, \lambda) d\lambda \quad (9)$$

$$= 1 + \langle A(a)B(c) \rangle_\sigma. \quad (10)$$

Noticing that $\langle A(a)B(b) \rangle_\pi = \langle A(b)B(a) \rangle_\pi$, we denote this quantity $(ab)_\pi$. Then the inequality reads

$$|(ab)_\sigma - (bc)_\sigma| \leq 1 + (ac)_\sigma. \quad (11)$$

□

Theorem 2. *The predictions of quantum mechanics cannot be explained by a local hidden variable theory.*

Proof. Consider a system of two spins. Let the preparation σ be the singlet state

$$|\sigma\rangle = \frac{1}{\sqrt{2}} (|0\rangle \otimes |1\rangle - |1\rangle \otimes |0\rangle). \quad (12)$$

Let the random variables $A_\pi(a)$ and $B_\pi(b)$ be the outcomes of measurements of $(\sigma \cdot a) \otimes \mathbb{1}$ and $\mathbb{1} \otimes (\sigma \cdot b)$, respectively, for unit vectors a and b . Then the correlations are

$$(ab)_\sigma = \langle A(a)B(b) \rangle_\sigma = \langle \sigma | (\sigma \cdot a) \otimes (\sigma \cdot b) | \sigma \rangle = -a \cdot b. \quad (13)$$

Then for any a , $(aa)_\sigma = -1$, as in the statement of Bell's theorem. If there is to be an explanatory local hidden variable theory, then Bell's inequality must always hold:

$$|-a \cdot b + b \cdot c| \leq 1 - a \cdot c. \quad (14)$$

Consider the following measurements:

$$a = (1, 0, 1)/\sqrt{2} \quad b = (0, 0, 1) \quad c = (1, 0, 0). \quad (15)$$

Then the inequality reads

$$\left| -1/\sqrt{2} \right| \leq 1 - 1/\sqrt{2} \longrightarrow 1 \leq \sqrt{2} - 1 \longrightarrow 2 \leq \sqrt{2}. \quad (16)$$

This is a contradiction. □